

BVD SCREENING TEST

Binocular Vision Dysfunction Questionnaire (BVDQ™)

Take the test
online!



Name _____ Date _____

Phone Number _____ Email _____

FOR AGES 14 AND OLDER

Directions: For each of the following questions, please check the answer that best describes your situation. If you wear glasses or contact lenses, answer the questions assuming that you are wearing them.

Always = every day
Occasionally = less than once per week

Frequently = at least once per week
Never = never

ALWAYS
FREQUENTLY
OCCASIONALLY
NEVER

1	Do you have headaches and/or facial pain?				
2	Do you have pain in your eyes with eye movement?				
3	Do you experience neck or shoulder discomfort?				
4	Do you have dizziness and/or light headedness?				
5	Do you experience dizziness, light headedness, or nausea while performing close-up activities (computer work, reading, writing, etc.)?				
6	Do you experience dizziness, light headedness or nausea while performing far-distance activities (driving, television, movies, etc.)?				
7	Do you experience dizziness, light headedness, or nausea when bending down and standing back up, or when getting up quickly from a seated position?				
8	Do you feel unsteady or drift to one side while walking?				
9	Do you feel overwhelmed or anxious while walking in a large department store (Target, Wal-Mart, Costco, etc.)?				
10	Do you feel overwhelmed or anxious when in a crowd?				
11	Does riding in a car make you feel dizzy or uncomfortable?				
12	Do you experience anxiety or nervousness because of your dizziness?				
13	Do you ever find yourself with your head tilted to one side?				
14	Do you experience poor depth perception or have difficulty estimating distances accurately?				
15	Do you experience double/overlapping/shadowed vision at far distances?				
16	Do you experience double/overlapping/shadowed vision at near distances?				
17	Do you experience glare or have sensitivity to bright lights?				
18	Do you close or cover one eye with near or far tasks?				
19	Do you skip lines or lose your place when you are reading? Do you use your finger, ruler or other guides to maintain your position on the page?				
20	Do you tire easily with close-up tasks (computer work, reading, writing)?				
21	Do you experience blurred vision with far-distance activities (driving, television, movies, chalkboard at school, etc.)?				
22	Do you experience blurred vision with close-up activities (computer work, reading, writing, etc.)?				
23	Do you blink to 'clear up' distant objects after working at a desk or working with close-up activities (computer work, reading, writing, etc.)?				
24	Do you experience words running together while reading?				
25	Do you experience difficulty with reading or reading comprehension?				
TOTALS					

Continue to the back side to complete and score the questionnaire →

	YES	NO
Have you ever been diagnosed with a traumatic brain injury (TBI)?		
Have you ever been diagnosed with a concussion?		
Have you ever been diagnosed with a lazy eye?		
Have you ever been diagnosed with a reading disability?		
Have you ever had an eye operation?		

On an average day, how much are you bothered by symptoms listed here? Rate each symptom from 0 -10 0 = None of that symptom 10 = Worst		None	Worst		None	Worst
	Dizziness	0	1 2 3 4 5 6 7 8 9 10	Neckache	0	1 2 3 4 5 6 7 8 9 10
	Nausea	0	1 2 3 4 5 6 7 8 9 10	Unsteady when walking	0	1 2 3 4 5 6 7 8 9 10
	Anxiety	0	1 2 3 4 5 6 7 8 9 10	Sensitivity to light	0	1 2 3 4 5 6 7 8 9 10
	Headache	0	1 2 3 4 5 6 7 8 9 10	Reading difficulty	0	1 2 3 4 5 6 7 8 9 10
				Sound sensitivity	0	1 2 3 4 5 6 7 8 9 10

This questionnaire is designed to identify individuals whose symptoms (ex. headache, dizziness, anxiety, etc.) may be due to vision misalignment.

How to score this questionnaire:
Add the scores for questions 1 - 25 to get a TOTAL score.

Always = _____ x 3 Frequently = _____ x 2 Occasionally = _____ x 1 Never = _____ x 0 **TOTAL** _____

If your Total Score is 15 or higher, schedule an appointment today.

Treatment is low cost and uses microprism lenses, and if indicated, noise cancelling devices. The average patient requires two appointments and two sets of lenses over 6-8 weeks, and can expect to experience an 80% reduction of symptoms.

Connect with a scheduler today:

SCAN TO CALL



CALL / TEXT

(904) 713-2020

DAU FAMILY EYE CARE
132 Everest Lane, Suite 5 Saint Johns, FL 32259